

# MATH 620: Variational Methods and Optimization I

Instructor: Prof. Wolfgang Bangerth  
Weber 214  
[bangerth@colostate.edu](mailto:bangerth@colostate.edu)

Lectures: Engineering E 206, Mondays/Wednesdays/Fridays, 12-12:50pm  
Office hours: Wednesdays, 1-2pm; or by appointment.

## Homework assignment 6 – due Friday 11/30/2018

**Problem 1 (A small variation for the Dirichlet problem).** In class, we have gone through the details of a proof for guaranteeing that a minimizer exists for the functional

$$I(u) = \int_{\Omega} |\nabla u|^2$$

over the (affine) space

$$X_g = \{u \in W^{1,2}(\Omega) : u|_{\partial\Omega} = g\}.$$

Among the other consequences of the theorem were that the (unique) minimizer  $\bar{u}$  had to satisfy the weak Euler-Lagrange equation

$$\int_{\Omega} \nabla \bar{u} \cdot \nabla \varphi = 0 \quad \forall \varphi \in X_0,$$

where  $X_0$  is the tangent space to  $X_g$  (i.e., consists of functions with zero boundary values), and that if  $\bar{u}$  happens to be smooth enough, that it then have to satisfy the partial differential equation

$$\begin{aligned} -\Delta \bar{u} &= 0 && \text{in } \Omega, \\ \bar{u} &= g && \text{on } \partial\Omega, \end{aligned}$$

i.e., it has to solve the Laplace equation.

Repeat some of the steps of the proof for the following variation:

$$I(u) = \int_{\Omega} |\nabla u|^2 - hu,$$

where  $h \in L^2(\Omega)$  is a given function. For simplicity take  $X_0 = W_0^{1,2}$  as the set to find a minimum over, i.e.,  $g = 0$ .

In particular, do the following:

- Repeat the first step of showing that a minimizer exists. Namely, we needed to show that for a minimizing sequence  $\{u_n\} \subset X_g$  so that  $I(u_n) \rightarrow m = \inf_{u \in X_g} I(u)$ , there exists an  $N$  and  $\gamma < \infty$  so that for all  $n \geq N$ , we have that  $\|u_n\|_{W^{1,2}} \leq \gamma$ .

$$\|u\|_{W^{1,2}} \leq \gamma.$$

The key to this was to show that

$$\|u\|_{W^{1,2}}^2 \leq c_1 I(u) + c_2.$$

If this is true, then we know – because  $u_n$  is a *minimizing sequence* – that there are  $N < \infty, |a| < \infty, b < \infty$  so that

$$I(u_n) \leq am + b$$

for all sufficiently large  $n \geq N$ . As a consequence, we know that after that point in the sequence,  $\|u\|_{W^{1,2}} \leq \sqrt{c_1(am+b) + c_2} = \gamma$  and the weak compactness of the ball of radius  $\gamma$  in  $W^{1,2}$  then guarantees that there is a weakly convergent subsequence.

Show a similar proof with the variation of the functional  $I(u)$  above.

- Show the weak Euler-Lagrange equation a minimizer has to satisfy.
- Show the strong Euler-Lagrange equation a minimizer has to satisfy if it is regular (smooth) enough.

(40 points)

The remainder of the homework is concerned with finding counter-examples for extensions of the general theorem we have mentioned in class. It read as follows:

**Theorem:** Let  $\Omega \subset \mathbb{R}^n$  be a bounded open set with a Lipschitz boundary. Let  $f \in C^0(\Omega \times \mathbb{R} \times \mathbb{R}^n)$ ,  $f = f(x, u, \xi)$  be a function that satisfies the following conditions:

- (i)  $\xi \mapsto f(x, u, \xi)$  is convex for all  $x \in \Omega, u \in \mathbb{R}$ ;
- (ii) there exist  $p > q \geq 1$  and  $\alpha_1 > 0, \alpha_2, \alpha_3 \in \mathbb{R}$  (i.e., they must be finite) so that

$$f(x, u, \xi) \geq \alpha_1 |\xi|^p + \alpha_2 |u|^q + \alpha_3$$

for all  $x \in \Omega, u \in \mathbb{R}, \xi \in \mathbb{R}^n$ .

Then the functional

$$I(u) = \int_{\Omega} f(x, u(x), \nabla u(x)) \, dx$$

has a minimizer  $\bar{u}$  in

$$X_g = \{u \in W^{1,p}(\Omega) : u|_{\partial\Omega} = g\},$$

where  $g$  is the restriction of some  $\tilde{g} \in W^{1,p}(\Omega)$  to  $\partial\Omega$ . (Or viewed differently,  $g$  are prescribed boundary values that are nice enough so that we can find an extension of  $g$  called  $\tilde{g}$  so that  $\tilde{g} \in W^{1,p}(\Omega)$  and so that  $\tilde{g}|_{\partial\Omega} = g$ .)

If, furthermore,

- (iii)  $f \in C^1$  and if there is a  $\beta \geq 0$  so that

$$\begin{aligned} |f_u(x, u, \xi)| &\leq \beta(1 + |u|^{p-1} + |\xi|^{p-1}), \\ |f_{\xi}(x, u, \xi)| &\leq \beta(1 + |u|^{p-1} + |\xi|^{p-1}), \end{aligned}$$

for all  $x \in \Omega, u \in \mathbb{R}, \xi \in \mathbb{R}^n$ ,

then  $\bar{u}$  satisfies the weak Euler-Lagrange equations

$$\int_{\Omega} (f_u(x, \bar{u}(x), \nabla \bar{u}(x))\varphi + f_{\xi}(x, \bar{u}(x), \nabla \bar{u}(x)) \cdot \nabla \varphi) \, dx = 0$$

for all  $\varphi \in X_0$ .

The theorem as stated seems to have a lot of restrictions, but it turns out that they all seem necessary since one can find counter-examples without too much trouble. The following exercises are therefore meant to probe the applicability of the theorem.

**Problem 2 (Application 1 of the general theorem).** Consider the function  $f(x, u, \xi) = \frac{1}{4}|\xi|^4 + g(x, u)$  where  $g \in C^{0,1}(\Omega \times \mathbb{R})$ . Show that the theorem applies. **(20 points)**

**Problem 3 (Application 2 of the general theorem).** Consider the function  $f(x, u, \xi) = \frac{1}{2}|\xi|^2 - \frac{1}{2}\lambda^2 u^2$  where  $\lambda$  is large – say, larger than the constant in the Poincaré inequality for functions in  $W_0^{1,2}(\Omega)$ . Show that the theorem does not apply by checking each condition individually. Then try to construct a sequence  $u_n$  so that  $I(u_n) \rightarrow -\infty$ , i.e., show that  $I(u)$  is not bounded from below on  $X_0 = W_0^{1,2}$ . For this part of the example, choose  $\Omega = (0, 1)$  and  $\lambda > \pi$ . **(20 points)**

**Problem 4 (Application 3 of the general theorem).** Consider the function  $f(x, u, \xi) = (|\xi|^2 - 1)^2$  on  $\Omega = (0, 1) \subset \mathbb{R}$  and with  $X_g = W_0^{1,4}(0, 1)$ . Show that the theorem does not apply by checking each condition individually.

Derive the weak and strong Euler-Lagrange equations for this case. Show that  $u = 0$  satisfies both of these equations; then show that it is not a minimizer of  $I(u)$ , for example by finding another function  $v \in X_g$  so that  $I(v) < I(u)$ . **(20 points)**