


The Mollifier Theorem

Definition of the Mollifier

The function

$$T(x) = \begin{cases} K \exp\left(\frac{-1}{1-|x|^2}\right) & \text{if } |x| < 1 \\ 0 & \text{if } |x| \geq 1 \end{cases}, \quad x \in \mathbb{R}^n$$

where the constant K is chosen such that $\int_{\mathbb{R}^n} T(x) dx = 1$, is a test function on \mathbb{R}^n . Note that $T(x)$ vanishes, together with all its derivatives as $|x| \rightarrow 1^-$, so $T(x)$ is infinitely differentiable and has compact support. The graph of $T(x)$ is sketched in the following figure.



The Mollifier Function

.For $n = 1$ and $\varepsilon > 0$, let

$$S_\varepsilon(x) = \frac{1}{\varepsilon} T\left(\frac{x}{\varepsilon}\right) \quad \text{and} \quad P_\varepsilon(x) = T\left(\frac{x}{\varepsilon}\right).$$

Then

$$\begin{aligned} S_\varepsilon(x) &\geq 0 & \text{and} & & P_\varepsilon(x) &\geq 0 & \text{for all } x \\ S_\varepsilon(x) &= 0 & \text{and} & & P_\varepsilon(x) &= 0 & \text{for } |x| > \varepsilon \\ \int_{\mathbb{R}} S_\varepsilon(x) dx &= 1 & \forall \varepsilon > 0, & & S_\varepsilon(0) &\rightarrow +\infty & \text{as } \varepsilon \rightarrow 0, \\ \int_{\mathbb{R}} P_\varepsilon(x) dx &\rightarrow 0 & \text{as } \varepsilon \rightarrow 0, & & P_\varepsilon(0) &= K/e & \forall \varepsilon > 0, \end{aligned}$$

Evidently, $S_\varepsilon(x)$ becomes thinner and higher as ε tends to zero but the area under the graph is constantly equal to one. On the other hand, $P_\varepsilon(x)$ has constant height but grows thinner as ε tends to zero. These test functions can be used as the "seeds" from which an infinite variety of other test functions can be constructed by using a technique called regularization which we will now describe.

For $n \geq 1$ we have

$$S_\varepsilon(x) = \frac{1}{\varepsilon^n} T\left(\frac{x}{\varepsilon}\right) \quad \text{and} \quad P_\varepsilon(x) = T\left(\frac{x}{\varepsilon}\right).$$

For U a bounded open set in R^n , and for $u \in L^1_{loc}(U)$, define for any $\varepsilon > 0$ and any $x \in U_\varepsilon = \{x \in U : \text{dist}(x, \partial U) > \varepsilon\}$,

$$J_\varepsilon u(x) = \int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y) u(y) dy \quad (1.1a)$$

$$= \int_{|z| \leq \varepsilon} S_\varepsilon(z) u(x-z) dz \quad (1.1b)$$

$$= \int_{|z| \leq 1} S_1(z) u(x-\varepsilon z) dz. \quad (1.1c)$$

We refer to $J_\varepsilon u(x)$ as the **mollified** $u(x)$. This mollified function, $J_\varepsilon u(x)$, is a smoothed version of the original function, $u(x)$.

Properties of the Mollifier

Note first that $J_\varepsilon u(x)$ is infinitely differentiable; i.e., for any $\varepsilon > 0$ and any $x \in U_\varepsilon$, it is clear from (1.1a) that

$$\frac{J_\varepsilon u(x + \varepsilon \vec{e}_i) - J_\varepsilon u(x)}{\varepsilon} = \int_{|x-y| \leq \varepsilon} \frac{[S_\varepsilon(x + \varepsilon \vec{e}_i - y) - S_\varepsilon(x - y)]}{\varepsilon} u(y) dy$$

i.e.,

$$\frac{J_\varepsilon u(x + \varepsilon \vec{e}_i) - J_\varepsilon u(x)}{\varepsilon} \rightarrow \int_{|x-y| \leq \varepsilon} \partial_{x_i} S_\varepsilon(x-y) u(y) dy \quad \text{as } \varepsilon \rightarrow 0.$$

Since $S_\varepsilon(x)$ is infinitely differentiable, it follows that $J_\varepsilon u(x)$ is infinitely differentiable on the open set U_ε .

It is evident from (1.1a) that for $1 \leq p < \infty$, $\varepsilon > 0$, and $x \in U_\varepsilon$,

$$J_\varepsilon u(x) = \int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y)^{1-1/p} S_\varepsilon(x-y)^{1/p} u(y) dy.$$

Then, using Holder's inequality, we get

$$|J_\varepsilon u(x)|^p = \left(\int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y) dy \right)^{p-1} \int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y) |u(y)|^p dy$$

and since $\int_R S_\varepsilon(x) dx = 1 \quad \forall \varepsilon > 0$,

$$\begin{aligned} \int_V |J_\varepsilon u(x)|^p dx &\leq \int_V \int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y) |u(y)|^p dy dx \\ &= \int_W |u(y)|^p \int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y) dx dy = \int_W |u(y)|^p dy \end{aligned}$$

for open sets $W = U_\varepsilon$, and $V = W_\varepsilon$. This result is just that assertion that

$$\|J_\varepsilon u\|_{L^p(V)} \leq \|u\|_{L^p(W)} \quad \text{for } V \subset\subset W \subset\subset U \quad (1.2)$$

Next, use (1.1c) to write

$$J_\varepsilon u(x) - u(x) = \int_{|z| \leq 1} S_1(z) [u(x-\varepsilon z) - u(z)] dz.$$

If the function $u = u(x)$ is, in fact, continuous on U , then this last result shows that

$$\max_V |J_\varepsilon u(x) - u(x)| \leq \max_V |[u(x-\varepsilon z) - u(z)]| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0; \quad (1.3)$$

i.e., $J_\varepsilon u(x)$ converges uniformly to $u(x)$ for $x \in \bar{V}$ when $u(x)$ is continuous on U .

For $u \in L^p_{loc}(U)$, $W = U_\varepsilon$, and arbitrary $\delta > 0$, use the fact that the continuous functions

are dense in $L_p(W)$ to choose $v \in C(W)$ such that

$$\|u - v\|_{L_p(W)} \leq \delta.$$

Then for $V = W_\varepsilon$,

$$\begin{aligned} \|J_\varepsilon u - u\|_{L_p(V)} &\leq \|J_\varepsilon u - J_\varepsilon v\|_{L_p(V)} + \|J_\varepsilon v - v\|_{L_p(V)} + \|v - u\|_{L_p(V)} \\ &\leq \|u - v\|_{L_p(W)} + \|J_\varepsilon v - v\|_{L_p(V)} + \|v - u\|_{L_p(W)} \leq 2\delta + \|J_\varepsilon v - v\|_{L_p(V)} \end{aligned}$$

It follows now from (1.3) that for $u \in L_{loc}^p(U)$,

$$\forall V \subset\subset U, \quad \|J_\varepsilon u - u\|_{L_p(V)} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0 \quad (1.4)$$

We can summarize these results in the following,

Theorem (Local Approximation) Suppose U is open and bounded in R^n , $1 \leq p < \infty$, and for $\varepsilon > 0$, let U_ε denote the subset $\{x \in U : \text{dist}(x, \partial U) > \varepsilon\}$.

(a) For every $\varepsilon > 0$, $u \in L_{loc}^p(U)$ implies $J_\varepsilon u \in C^\infty(U_\varepsilon)$

(b) (i) $u \in C(U)$ implies u_ε converges to u uniformly on compact subsets of U ; i.e.,

$$\|J_\varepsilon u - u\|_{C(\bar{V})} = \max_{\bar{V}} |J_\varepsilon u(x) - u(x)| \rightarrow 0 \text{ for all } V \subset\subset U$$

(ii) u_ε converges to u in $L_{loc}^p(U)$; i.e., $u \in L_{loc}^p(U)$ implies that for all $V \subset\subset W \subset\subset U$,

$$\|J_\varepsilon u\|_{L^p(V)} \leq \|u\|_{L^p(W)} \quad \text{and} \quad \|J_\varepsilon u - u\|_{L^p(V)} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0$$

(c) u_ε converges to u in $W_{loc}^{k,p}(U)$;

Result (c) follows from (b) by induction.

Corollary (Global Approximation) Suppose U has a smooth boundary, and $1 \leq p < \infty$.

(a) For every $\varepsilon > 0$, $u \in L^p(U)$ implies $J_\varepsilon u \in C^\infty(U) \cap L^p(U)$.

(b) $u \in L^p(U)$ implies that

$$\|J_\varepsilon u\|_{L^p(U)} \leq \|u\|_{L^p(U)} \quad \text{and} \quad \|J_\varepsilon u - u\|_{L^p(U)} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0$$

(c) $u \in W^{k,p}(U)$ implies that there exists functions $\{\phi_m\} \in C^\infty(U) \cap W^{k,p}(U)$ such that

$$\|\phi_m - u\|_{k,p} \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

The proof of the corollary makes use of a partition of unity (see theorem 2 pg 251 in Evans).

Weak Equals Strong

For U a bounded open set in R^n , we define $v = v(x)$ to be the **weak derivative** of order α , of $u = u(x)$, $x \in U$ if

$$\int_U u(x) \partial^\alpha \phi(x) dx = (-1)^{|\alpha|} \int_U v(x) \phi(x) dx \text{ for all } \phi \in C_c^\infty(U)$$

Similarly, we define $v = v(x)$ to be the **strong L_p -derivative** of order α , of $u = u(x)$, $x \in U$ if

for any $V \subset\subset U$, there exists a sequence $\{\phi_n\} \in C_c^\infty(U)$ such that

$$\int_V |\phi_n - u|^p dx \rightarrow 0 \quad \text{and} \quad \int_V |\partial^\alpha \phi_n - v|^p dx \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Using mollifiers, we can show that these two notions are equivalent.

Suppose first that $v = v(x)$ is the weak derivative of order α , of $u = u(x)$. Then, since $S_\varepsilon \in C_c^\infty(U)$,

$$\begin{aligned} \partial^\alpha J_\varepsilon u(x) &= \int_{|x-y| \leq \varepsilon} \partial_x^\alpha S_\varepsilon(x-y) u(y) dy = (-1)^{|\alpha|} \int_{|x-y| \leq \varepsilon} \partial_y^\alpha S_\varepsilon(x-y) u(y) dy \\ &= \int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y) v(y) dy = J_\varepsilon v(x) \quad (\text{by definition of weak derivative}) \end{aligned}$$

Now apply (1.4) to write

$$\int_V |J_\varepsilon u - u|^p dx \rightarrow 0 \quad \text{and} \quad \int_V |\partial^\alpha J_\varepsilon u - v|^p dx = \int_V |J_\varepsilon v - v|^p dx \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Thus every weak derivative is a strong L_p -derivative.

Conversely, suppose $v = v(x)$ is the strong L_p -derivative of order α , of $u = u(x)$ with

$$\int_V |\phi_n - u|^p dx \rightarrow 0 \quad \text{and} \quad \int_V |\partial^\alpha \phi_n - v|^p dx \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

for arbitrary $V \subset\subset U$, and $\{\phi_n\} \in C_c^\infty(U)$. Then for any $\psi \in C_c^\infty(U)$,

$$\begin{aligned} \int_V (u - \phi_n) \partial^\alpha \psi dx &= \int_V u \partial^\alpha \psi dx - \int_V \phi_n \partial^\alpha \psi dx \\ &= \int_V u \partial^\alpha \psi dx - (-1)^{|\alpha|} \int_V \partial^\alpha \phi_n \psi dx \\ &= \int_V u \partial^\alpha \psi dx - (-1)^{|\alpha|} \int_V v \psi dx + (-1)^{|\alpha|} \int_V (v - \partial^\alpha \phi_n) \psi dx \end{aligned}$$

Then it follows that

$$\left| \int_V u \partial^\alpha \psi dx - (-1)^{|\alpha|} \int_V v \psi dx \right| \leq C_1 \int_V |\phi_n - u|^p dx + C_2 \int_V |\partial^\alpha \phi_n - v|^p dx$$

which implies that every strong L_p -derivative is a weak derivative.

Weyl's Lemma

Weyl's lemma is a famous result that asserts that for U a bounded open set in R^n , if $u = u(x)$ is harmonic in U , (i.e., $u \in C^2(U)$ and $\nabla^2 u(x) = 0$, $x \in U$) then $u(x)$ is infinitely differentiable in U .

To see why this result is true, recall that every harmonic function has the mean value property. That is,

$$\forall x \in U_\varepsilon, r < \varepsilon, \quad u(x) = \int_{\partial B_r(x)} u(y) d\hat{S}(y) = \frac{1}{nr^{n-1}A_n} \int_{\partial B_r(x)} u(y) dS(y).$$

Then

$$\begin{aligned} J_\varepsilon u(x) &= \int_{|x-y| \leq \varepsilon} S_\varepsilon(x-y) u(y) dy = \frac{1}{\varepsilon^n} \int_{|x-y| \leq \varepsilon} T\left(\frac{x-y}{\varepsilon}\right) u(y) dy \\ &= \frac{1}{\varepsilon^n} \int_0^\varepsilon T\left(\frac{r}{\varepsilon}\right) \int_{\partial B_r(x)} u(y) dS(y) dr = u(x) \int_0^\varepsilon \frac{nA_n}{\varepsilon^n} T\left(\frac{r}{\varepsilon}\right) r^{n-1} dr \\ &= u(x) \int_{B_\varepsilon(0)} S_\varepsilon(y) dy = u(x). \end{aligned}$$

But this says that $\forall \varepsilon > 0$, $\forall x \in U_\varepsilon$, $J_\varepsilon u(x) = u(x)$. Since $J_\varepsilon u(x)$ is infinitely differentiable on U , it follows that $u(x)$ is infinitely differentiable on U although u need not even be continuous on the closure, \bar{U} .