


Borsuk-Ulam Theorems and Vietoris-Rips complexes

Joint with Johnathan Bush and Florian Frick
Mathematika 2020




Emphasis: Connections to ...

(I) Borsuk-Ulam Theorem



Given $f: S^n \rightarrow \mathbb{R}^n$, $\exists x \in S^n$ with

(II) What about $S^n \rightarrow \mathbb{R}^k$ with $k \leq n$?




Gromov's Waist Inequality For $f: S^n \rightarrow \mathbb{R}^k$ with $k \leq n$,
 $\exists y \in \mathbb{R}^k$ with $\text{Vol}_{n-k}(f^{-1}(y)) \geq \text{Vol}_{n-k}(\text{Sphere})$.

Proof Almgren: 100 pages of geometric measure theory
Gromov: The filling radius, or characteristic classes.
Lim, Mémoli, Okutan connect to Vietoris-Rips complexes

Remark Implies invariance of dimension:


(III) What about $f: S^1 \rightarrow \mathbb{R}^k$ with $k \geq n$?



S^1 with path-length metric, unit circumference.

Theorem (A, Bush, Frick) For $f: S^1 \rightarrow \mathbb{R}^{2k+1}$,
 $\exists X \subset S^1$ of diameter at most $\frac{k}{2k+1}$
such that $\text{conv}() \cap \text{conv}() \neq \emptyset$.

$$r = \frac{1}{3} \text{ (triangle)} \quad r = \frac{2}{5} \text{ (pentagon)} \quad r = \frac{3}{7} \text{ (heptagon)}$$



Sharpness of diameter bound

$$S^1 \xrightarrow{\quad} \mathbb{R}^{2k} \subseteq \mathbb{R}^{2k+1}$$




$$\theta \mapsto (\cos \theta, \sin \theta, \quad)$$

Proof


$$S^1 \xrightarrow{s} \mathbb{R}^{2k+1} \text{ induces}$$

$$\text{VR}(S^1; \frac{k}{2k+1}) \xrightarrow{s} \mathbb{R}^{2k+1}$$

Vietoris-Rips simplicial complex with vertex set S^1 containing all simplices of diameter




Theorem (A, Bush, Frick) For $f: S^n \rightarrow \mathbb{R}^{n+2}$,
 $\exists X \subset S^n$ of diameter at most r_n
such that $\text{conv}(f(X)) \cap \text{conv}(f(-X)) \neq \emptyset$.




Proof

$$S^n \xrightarrow{f} \mathbb{R}^{n+2} \quad \text{induces}$$

$$(n+1)\text{-connected } \text{VR}(S^n; r_n) \xrightarrow{f} \mathbb{R}^{n+2}$$



Remark Lovász' strongly self-dual polytopes.



Remark Michael Crabb uses characteristic classes
to get extensions

$$S^{2r+t} \longrightarrow$$

Remark Versions of the ham sandwich theorem
with more "fixings" than the dimension!

Remark Quantitative topology, esp.

Isosystolic Inequality:

$$\text{sys}(M) \leq C \text{ vol}(M)^{\frac{1}{n}}$$

1983 Mikhael Gromov

1983-1991 Mikhail Katz

2020 Lim, Memoli, Okutan

2017 Adamszek, A


2018 Adamszek, A, Frick

2020 A, Bush, Frick

(IV) Orbitopes and Schur polynomials

Defⁿ The Barvinok-Novik orbitope $B_{2k} \subseteq \mathbb{R}^{2k}$ is the convex hull of the curve $(\cos \theta, \sin \theta, \cos 3\theta, \sin 3\theta, \cos 5\theta, \sin 5\theta, \dots, \cos(2k-1)\theta, \sin(2k-1)\theta)$.

Its faces are known only for $k=1, 2$.



Defⁿ The Carathéodory orbitope $C_{2k} \subseteq \mathbb{R}^{2k}$ is the convex hull of the curve $(\cos \theta, \sin \theta, \cos 2\theta, \sin 2\theta, \dots, \cos k\theta, \sin k\theta)$.

Its faces are known for all k .



	Odd Barvinok-Novik	Not odd Carathéodory
(scale) ↓ What diameter hits origin?	$\cos t_1, \cos t_2, \cos t_3, \cos t_4$ $\sin t_1, \sin t_2, \sin t_3, \sin t_4$ $\cos 3t_1, \cos 3t_2, \cos 3t_3, \cos 3t_4$ $\sin 3t_1, \sin 3t_2, \sin 3t_3, \sin 3t_4$	$\cos t_1, \cos t_2, \cos t_3, \cos t_4$ $\sin t_1, \sin t_2, \sin t_3, \sin t_4$ $\cos 2t_1, \cos 2t_2, \cos 2t_3, \cos 2t_4$ $\sin 2t_1, \sin 2t_2, \sin 2t_3, \sin 2t_4$
What are the orbitope's faces?	1 1 1 1 1 $\cos t_1, \cos t_2, \cos t_3, \cos t_4, \cos t_5$ $\sin t_1, \sin t_2, \sin t_3, \sin t_4, \sin t_5$ $\cos 3t_1, \cos 3t_2, \cos 3t_3, \cos 3t_4, \cos 3t_5$ $\sin 3t_1, \sin 3t_2, \sin 3t_3, \sin 3t_4, \sin 3t_5$	1 1 1 1 1 $\cos t_1, \cos t_2, \cos t_3, \cos t_4, \cos t_5$ $\sin t_1, \sin t_2, \sin t_3, \sin t_4, \sin t_5$ $\cos 2t_1, \cos 2t_2, \cos 2t_3, \cos 2t_4, \cos 2t_5$ $\sin 2t_1, \sin 2t_2, \sin 2t_3, \sin 2t_4, \sin 2t_5$

Changing to exponentials gives

- A Vandermonde matrix, whose determinant is easy to factor, answering the question.
- A generalized Vandermonde matrix, whose determinant contains a Schur polynomial in its factorization, whose sign is hard to analyze.

$$\text{Top left: } \det = C \prod_{i < j} \sin(t_j - t_i)$$

$$\text{Bottom right: } \det = C \prod_{i < j} \sin\left(\frac{t_j - t_i}{2}\right)$$

$$\text{Bottom left: } \det = C \left(\prod_{1 \leq i < j \leq 5} \sin\left(\frac{t_j - t_i}{2}\right) \right) \left(2 + \sum_{1 \leq i < j \leq 5} \cos(t_j - t_i) \right)$$

Vandermonde Schur