

MATH 609-602: Numerical Methods

Lecturer: Prof. Wolfgang Bangerth
Blocker Bldg., Room 507D
(979) 845 6393
bangerth@math.tamu.edu

Teaching Assistant: Seungil Kim
Blocker Bldg., Room 507A
(979) 862 3259
sgkim@math.tamu.edu

Homework assignment 9 – due Tuesday 11/8/2005

Problem 1 (Finite difference approximation of the derivative). Take the function defined by

$$f(x) = \begin{cases} \frac{1}{2}x^3 + x^2 & \text{for } x < 0 \\ x^3 & \text{for } x \geq 0. \end{cases}$$

Compute a finite difference approximation to $f'(x_0)$ at $x_0 = 1$ with both the one-sided and the symmetric two-sided formula. Use step sizes $h = 1, \frac{1}{2}, \frac{1}{4}, \dots, \frac{1}{64}$. Determine experimentally the convergence orders you observe as $h \rightarrow 0$.

Repeat these computations for $x_0 = 0$. What convergence orders do you observe? Why? **(4 points)**

Problem 2 (Derivatives of an implicit function). Let $f(x)$ be defined implicitly as follows: for every $x > 0$, $f(x)$ is that value y for which

$$ye^y = x. \tag{1}$$

In other words, every time one wants to evaluate $f(x)$ for a particular value x , one has to solve equation (1) for y . This can be done using Newton's method, for example, or any of the other root finding algorithms we had in class applied to the function $g(y) = ye^y - x$.

Plot $f(x)$ for $0 \leq x \leq 10$. Compute an accurate approximation to $f'(2)$.

(4 points)

Problem 3 (Integration of an implicit function). Let $f(x)$ be defined as in Problem 2. Compute

$$\int_0^{10} f(x) dx$$

using both the box rule as well as the trapezoidal rule for step sizes $h = 1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots, \frac{1}{32}$. Determine the order of convergence for both methods.

(4 points)

Problem 4 (A proof). In last week's homework, you were asked to find the l_∞ best approximating linear function to 10 data points. Let's simplify the situation a little bit: assume we had only wanted to find a *constant* best approximation, i.e. a function $p_0(x) = c_0$, to all these data points. Then, this involves finding the coefficient c_0^* for which the function

$$e(c_0) = \max_{1 \leq i \leq N} |c_0 - y_i|$$

takes on its minimum. If one wants to phrase this differently, one could also say that we are looking for the optimal coefficient c_0^* for which

$$e(c_0^*) = \min_{c_0} \max_{1 \leq i \leq N} |c_0 - y_i|.$$

One could wonder if there is indeed only a single such value c_0 , or if it may be possible to have a number of *different* values for c_0 for which the corresponding functions $p_0(x) = c_0$ are all best approximations to the data points.

Prove that the function $e(c_0)$ defined above has only a single minimum and that consequently there is exactly one, well-defined best l_∞ approximation among the constant functions. (Hint: Try your hands first on the case where there are only two data points, i.e. $e(c_0) = \max\{|c_0 - y_1|, |c_0 - y_2|\}$ and then generalize to N data points.)

Comment on what happens if you were looking for linear approximations $p_1(x) = c_0 + c_1x$ with corresponding error function

$$e(c_0, c_1) = \max_{1 \leq i \leq N} |c_0 + c_1x_i - y_i|.$$

Does this two-dimensional function have a single, unique minimum as well?

(4 points)