

Statement of Recent Work


Henry Adams (adams@math.colostate.edu)

Given a set of points X sampled from a metric space M , what information can one recover about the underlying space M ? One approach is to build a *Vietoris–Rips simplicial complex* $\text{VR}(X; r)$ on top of the sampling X ; this is a geometric construction which depends on the choice of a scale parameter $r > 0$. In nice cases the Vietoris–Rips complex of the sampling X will recover the unknown space M up to homotopy type. Vietoris–Rips complexes were first invented for use in geometric group theory and metric cohomology [12, 22], but within the last 20 years they have emerged as an important tool in applied and computational topology, including persistent homology [13, 15]. In particular, data science practitioners currently use software packages to compute Vietoris–Rips complexes of datasets X as a way to approximate the shape of the dataset. However, in order to interpret their computations, there is a need to mathematically better understand the limiting object (the Vietoris–Rips complex of the underlying shape M). My recent work has extended the theory of Vietoris–Rips complexes (§1) at larger scales, inspired a new definition of these thickenings allowing intuitive proofs of old and new results (§2), and connected the mathematics to interdisciplinary research in machine learning, sensor networks, and high-dimensional data analysis (§3).

§1. Vietoris–Rips simplicial complexes at larger scale parameters.

Definition. *Given a metric space X and scale parameter $r \geq 0$, the Vietoris–Rips simplicial complex $\text{VR}(X; r)$ contains a finite subset $\sigma \subseteq X$ as a simplex if its diameter is at most r .*

Since we do not know a priori how to choose the scale r , the idea of persistent homology is to compute the homology of the Vietoris–Rips complex of metric space X over a large range of scale parameters r , and to trust those topological features which persist. The motivation for using Vietoris–Rips complexes are Hausmann’s and Latschev’s reconstruction results [19, 21]: for M a Riemannian manifold, scale r sufficiently small, and sample $X \subseteq M$ sufficiently dense, we have a homotopy equivalence $\text{VR}(X; r) \simeq M$. But as the key idea of persistence is to allow scale r to vary, the assumption that scale r is kept sufficiently small does not hold in practice.



Theorem 1 ([1]). *As the scale r increases, the Vietoris–Rips complexes of the circle are homotopy equivalent to the circle, the 3-sphere, the 5-sphere, the 7-sphere, . . .*

This is the first connected non-contractible Riemannian manifold for which the homotopy types of the Vietoris–Rips complex are known at all choices of scale. We extend our result to classify the homotopy types of Vietoris–Rips complexes of ellipses, regular polygons, n -dimensional tori, random subsets from the circle, and metric gluings [3, 4, 5, 6]. Our work has also been extended in [17, 27, 26] in order to identify the 1-dimensional persistent homology of metric graphs and geodesic spaces, and in [28] for applications to geometric group theory. However, in general little is known about Vietoris–Rips complexes at larger scales, though they arise naturally in applications of persistent homology.

Question. *What are the homotopy types of Vietoris–Rips spaces of Riemannian manifolds?*

The above question is fundamental for applications of persistent homology to data analysis, since as finite subset $X \subseteq M$ gets denser and denser, the persistent homology of the Vietoris–Rips complex of X converges to the (unknown) persistent homology of the Vietoris–Rips complex of manifold M [14]. In the following section, we give a partial answer for n -spheres.

§2. From Vietoris–Rips complexes to metric thickenings. Though we started with a metric space X , if X is not discrete then the Vietoris–Rips complex $\text{VR}(X; r)$ is not a metric space. Indeed, if $\text{VR}(X; r)$ is not locally finite then it is not metrizable, meaning it is *impossible* to equip $\text{VR}(X; r)$ with a metric without changing the homeomorphism type. A related problem (especially for proofs) is that for X not discrete, the inclusion $X \hookrightarrow \text{VR}(X; r)$ is not continuous. In [2] we address these issues. We use optimal transport (i.e. a Wasserstein or Kantorovich metric [16, 18, 25]) to build a metric space $\text{VR}^m(X; r)$ — the *Vietoris–Rips thickening* — for all $r > 0$.

There is strong evidence that when X is not a discrete metric space, then the metric thickening $\text{VR}^m(X; r)$ is a more natural object than the simplicial complex $\text{VR}(X; r)$. Consider for example Hausmann’s theorem [19], which states that if M is a compact Riemannian manifold, then for small enough scale parameters $r > 0$, the complex $\text{VR}(M; r)$ is homotopy equivalent to M . Though this theorem is foundational for the subject, its proof is quite complicated. Indeed, Hausmann’s homotopy equivalence $T: \text{VR}(M; r) \rightarrow M$ is *extremely non-canonical*; it depends on the arbitrary choice of a fixed total ordering of all of the points in manifold M , and different choices of total orderings will produce different maps $T: \text{VR}(M; r) \rightarrow M$. Furthermore, since the manifold M is not a discrete metric space, the inclusion $M \hookrightarrow \text{VR}(M; r)$ is not continuous and hence cannot be a homotopy inverse for T . Indeed, Hausmann does not write down an explicit homotopy inverse for T . By contrast, when using metric Vietoris–Rips thickenings instead of simplicial complexes, we give a natural proof of a metric analogue of Hausmann’s theorem.

Theorem 2 ([2]). *For M a Riemannian manifold of bounded curvature, the Vietoris–Rips metric thickening $\text{VR}^m(M; r)$ is homotopy equivalent to M for r sufficiently small.*

In our proof we use Karcher means [20] to define a map $g: \text{VR}^m(M; r) \rightarrow M$ which is a homotopy equivalence for r sufficiently small. Key features of our proof are that we give a canonical choice for map g , which has the (now continuous) inclusion $\iota: M \hookrightarrow \text{VR}^m(M; r)$ as a homotopy inverse, as proven by writing down a simple linear homotopy from the composition $\iota \circ g$ to the identity map.

Using metric thickenings, in [2] we are able to identify the first new homotopy type in the Vietoris–Rips thickening of the n -sphere, namely $\text{VR}^m(S^n; r) \simeq \Sigma^{n+1} \frac{\text{SO}(n+1)}{A_{n+2}}$, with alternating group A_{n+2} arising as the rotational symmetries of an inscribed regular $(n+1)$ -simplex. Extensions to larger scale parameters will relate to Lovász’ strongly self-dual polytopes [23].

In future work, I am developing a Morse theory for Vietoris–Rips thickenings: one will be able to understand the homotopy types of Vietoris–Rips thickenings of nice metric space at all scales without having to invent new tools for each new space.

§3. Interdisciplinary applications. I have worked on applications of Vietoris–Rips complexes to machine learning [11], sensor networks [10], and high-dimensional data analysis [7, 8, 9]. I am also a coauthor of the tutorial for Javaplex [24], a software package for persistent homology, which contains exercises, solutions, and examples on real life data sets. I help advertise and share my broader research area in my role as the Director of the Applied Algebraic Topology Research Network, with over 400 members. Our research network hosts an online research seminar, with 85 hour-long research seminar recordings posted to our YouTube Channel, which has over 300 subscribers and averages 2.5 hours watched per day.

REFERENCES

- [1] Michał Adamaszek and Henry Adams. The Vietoris–Rips complexes of a circle. *Pacific Journal of Mathematics*, 290:1–40, 2017.
- [2] Michał Adamaszek, Henry Adams, and Florian Frick. Metric reconstruction via optimal transport. *SIAM Journal on Applied Algebra and Geometry*, 2(4):597–619, 2018.
- [3] Michał Adamaszek, Henry Adams, Florian Frick, Chris Peterson, and Corrine Previte-Johnson. Nerve complexes of circular arcs. *Discrete & Computational Geometry*, 56:251–273, 2016.
- [4] Michał Adamaszek, Henry Adams, Ellen Gasparovic, Maria Gommel, et al. Vietoris–Rips and Čech complexes of metric gluings. *Proc. 34th Symposium on Computational Geometry*, pages 3:1–3:15, 2018.
- [5] Michał Adamaszek, Henry Adams, and Francis Motta. Random cyclic dynamical systems. *Advances in Applied Mathematics*, 83:1–23, 2017.
- [6] Michał Adamaszek, Henry Adams, and Samadwara Reddy. On Vietoris–Rips complexes of ellipses. *Journal of Topology and Analysis*, pages 1–30, 2017.
- [7] Henry Adams, Manuchehr Aminian, Elin Farnell, Michael Kirby, Chris Peterson, Joshua Mirth, et al. A fractal dimension for measures via persistent homology. *arXiv preprint arXiv:1808.01079*, 2018.
- [8] Henry Adams, Johnathan Bush, Brittany Carr, Lara Kassab, and Joshua Mirth. On the nonlinear statistics of optical flow. *Proc. Computational Topology in Image Context*, LNCS 11382:151–165, 2019.
- [9] Henry Adams and Gunnar Carlsson. On the nonlinear statistics of range image patches. *SIAM Journal on Imaging Sciences*, 2(1):110–117, 2009.
- [10] Henry Adams and Gunnar Carlsson. Evasion paths in mobile sensor networks. *International Journal of Robotics Research*, 34(1):90–104, 2015.
- [11] Henry Adams, Sofya Chepushtanova, Tegan Emerson, Eric Hanson, et al. Persistence images: A vector representation of persistent homology. *Journal of Machine Learning Research*, 18(8):1–35, 2017.
- [12] Martin R Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319. Springer Science & Business Media, 2011.
- [13] Gunnar Carlsson. Topology and data. *Bull. of the American Mathematical Soc.*, 46(2):255–308, 2009.
- [14] Frédéric Chazal, Vin de Silva, and Steve Oudot. Persistence stability for geometric complexes. *Geometriae Dedicata*, pages 1–22, 2013.
- [15] H. Edelsbrunner and J. Harer. *Computational Topology: An Introduction*. AMS, Providence, 2010.
- [16] David A Edwards. On the Kantorovich–Rubinstein theorem. *Exp. Mathematicae*, 29(4):387–398, 2011.
- [17] Ellen Gasparovic et al. A complete characterization of the one-dimensional intrinsic Čech persistence diagrams for metric graphs. In *Research in Computational Topology*, pages 33–56. Springer, 2018.
- [18] Mikhail Gromov. Geometric group theory, volume 2: Asymptotic invariants of infinite groups. *London Mathematical Society Lecture Notes*, 182:1–295, 1993.
- [19] Jean-Claude Hausmann. On the Vietoris–Rips complexes and a cohomology theory for metric spaces. *Annals of Mathematics Studies*, 138:175–188, 1995.
- [20] Hermann Karcher. Riemannian center of mass and mollifier smoothing. *Communications on pure and applied mathematics*, 30(5):509–541, 1977.
- [21] Janko Latschev. Vietoris–Rips complexes of metric spaces near a closed Riemannian manifold. *Archiv der Mathematik*, 77(6):522–528, 2001.
- [22] Solomon Lefschetz. *Algebraic topology*, volume 27. American Mathematical Society, 1942.
- [23] László Lovász. Self-dual polytopes and the chromatic number of distance graphs on the sphere. *Acta Scientiarum Mathematicarum*, 45(1-4):317–323, 1983.
- [24] Andrew Tausz, Mikael Vejdemo-Johansson, and Henry Adams. Javaplex: A research software package for persistent (co)homology. In *International Congress on Mathematical Software*, pages 129–136, 2014.
- [25] Cédric Villani. *Topics in optimal transportation*. Number 58. American Mathematical Soc., 2003.
- [26] Žiga Virk. 1-dimensional intrinsic persistence of geodesic spaces. *Topology and Analysis*, 2019.
- [27] Žiga Virk. Approximations of 1-dimensional intrinsic persistence of geodesic spaces and their stability. *Revista Matemática Complutense*, 32:195–213, 2019.
- [28] Matthew C. B. Zaremsky. Bestvina–Brady discrete Morse theory and Vietoris–Rips complexes. *arXiv preprint arXiv:1812.10976*, 2018.