An Introduction to Applied Topology

Henry Adams
Colorado State University

An Introduction to Applied Topology

Part I: Topology applied to data analysis

Part II: Topology applied to sensor networks

Datasets have shapes

Example: Diabetes study
145 points in 5-dimensional space



An attempt to define the nature of chemical diabetes using a multidimensional analysis by G. M. Reaven and R. G. Miller, 1979

Datasets have shapes

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space



Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Datasets have shapes

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space



Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Datasets have shapes

Topology studies shapes

A donut and coffee mug are “homotopy equivalent”, and considered to be the same shape. You can bend and stretch (but not tear) one to get the other.



Topology studies shapes



Topological Analysis of Population Activity in Visual Cortex by Singh, Memoli, Ishkhanov, Sapiro, Carlsson, and Ringach, 2008.

Topology studies shapes

Torus



Topology studies shapes

Klein bottle



Topology studies shapes

Klein bottle





Image credit: https://plus.maths.org/content/imaging-maths-inside-klein-bottle

Topology studies shapes

Klein bottle







Image credit: https://plus.maths.org/content/imaging-maths-inside-klein-bottle

Homology

- i-dimensional homology H_i "counts the number of i-dimensional holes"
- i-dimensional homology H_i actually has the structure of a vector space!



0-dimensional homology H_0: rank 6

1-dimensional homology H_1: rank 0



0-dimensional homology H_0: rank 1

1-dimensional homology H_1: rank 3



0-dimensional homology H_0: rank 1

1-dimensional homology H_1: rank 6

Homology

- i-dimensional homology "counts the number of i-dimensional holes"
- i-dimensional homology actually has the structure of a vector space!



0-dimensional homology H_0: rank 1

1-dimensional homology H_1: rank 0

2-dimensional homology H_2: rank 1



0-dimensional homology H_0: rank 1

1-dimensional homology H_1: rank 2

2-dimensional homology H_2: rank 1



Be careful! (Same as torus over $\mathbb{Z}/2\mathbb{Z}$)

Image credit: https://plus.maths.org/content/imaging-maths-inside-klein-bottle

Topology studies shapes

What shape is this?



















Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\check{\text{C}}\text{ech}(X; r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.



Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\check{\text{C}}\text{ech}(X; r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.



Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\check{\text{C}}\text{ech}(X; r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.



Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\text{Čech}(X; r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.



Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\check{\text{C}}\text{ech}(X; r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.



Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\text{Čech}(X; r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.



Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\check{\text{C}}\text{ech}(X; r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.



<u>Nerve Lemma.</u> $\check{\mathrm{C}}\mathrm{ech}(X;r) \simeq$ union of balls

Definition

For a data set $X \subseteq \mathbb{R}^n$ and scale $r \geq 0$, the *Čech simplicial complex* $\check{\mathrm{C}}\mathrm{ech}(X;r)$ has

- vertex set X
- finite simplex $\{x_0, x_1, \ldots, x_k\}$ when $\cap_{i=0}^{k} B(x_i, r) \neq \emptyset$.

Persistent homology





- Input: Increasing spaces. Output: barcode.
- Significant features persist.
- Cubic computation time in the number of simplices.

Persistent homology



- Input: Increasing spaces. Output: barcode.
- Significant features persist.
- Cubic computation time in the number of simplices.

Persistent homology



- Input: Increasing spaces. Output: barcode.
- Significant features persist.
- Cubic computation time in the number of simplices.

Persistent homology



- Input: Increasing spaces. Output: barcode.
- Significant features persist.
- Cubic computation time in the number of simplices.

Persistent homology

Sublevelset persistence





Analysis of Kolmogorov flow and Rayleigh–Bénard convection using persistent homology by Miroslav Kramár, Rachel Levanger, Jeffrey Tithof, Balachandra Suri, Mu Xu, Mark Paul, Michael F Schatz, Konstantin Mischaikow

- Input: Increasing spaces. Output: barcode.
- Significant features persist.
- Cubic computation time in the number of simplices.

Persistent homology

Sublevelset persistence





Analysis of Kolmogorov flow and Rayleigh–Bénard convection using persistent homology by Miroslav Kramár, Rachel Levanger, Jeffrey Tithof, Balachandra Suri, Mu Xu, Mark Paul, Michael F Schatz, Konstantin Mischaikow

- Input: Increasing spaces. Output: barcode.
- Significant features persist.
- Cubic computation time in the number of simplices.

Persistent homology applied to data

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space





Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Persistent homology applied to data

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space





Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Persistent homology applied to data



Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Persistent homology applied to data

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space



Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Persistent homology applied to data

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space



Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Persistent homology applied to data

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space



Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Persistent homology applied to data

Example: Cyclo-Octane (C_8H_{16}) data

1,000,000+ points in 24-dimensional space





Non-Manifold Surface Reconstruction from High Dimensional Point Cloud Data
by Shawn Martin and Jean-Paul Watson, 2010.

Persistent homology applied to data

Example: Equilateral pentagons in the plane





Image credit: Clayton Shonkwiler



Persistent homology applied to data





Image credit: Clayton Shonkwiler



Persistent homology applied to data

- <u>Stability Theorem.</u>

  If X and Y are metric spaces, then

$$d_b(\mathrm{PH}(\check{\mathrm{C}}\mathrm{ech}(X)), \mathrm{PH}(\check{\mathrm{C}}\mathrm{ech}(Y))) \leq 2d_{\mathrm{GH}}(X, Y)$$



Topology applied to image data



Persistent homology applied to data

The receptive fields of cells in our primary visual cortex (V1) are related to the statistics natural images.

Independent component filters of natural images compared with simple cells in primary visual cortex by JH van Hateren and A van der Schaaf, 1997

Persistent homology applied to data

3x3 *high-contrast* patches from images

Points in 9-dimensional space, normalized to have average color gray and contrast norm one (on 7-sphere).



On the Local Behavior of Spaces of Natural Images by Gunnar Carlsson, Tigran Ishkhanov, Vin de Silva, and Afra Zomorodian, 2008.

Persistent homology applied to data

1. Densest patches according to a global estimate



Persistent homology applied to data

1. Densest patches according to a global estimate





Interpretation: nature prefers linearity

Persistent homology applied to data

2. Densest patches according to an intermediate estimate



Persistent homology applied to data

2. Densest patches according to an intermediate estimate





Interpretation: nature prefers horizontal and vertical directions

Persistent homology applied to data

3. Densest patches according to a local estimate



Persistent homology applied to data

3. Densest patches according to a local estimate



Persistent homology applied to data

3. Densest patches according to a local estimate



Persistent homology applied to data

3. Densest patches according to a local estimate





Image credit: https://plus.maths.org/content/imaging-maths-inside-klein-bottle

Interpretation: nature prefers linear and quadratic patches at all angles

Why is applied topology popular when few datasets have Klein bottles?

- Many datasets have clusters & flares (as in the diabetes example)
- Motivates interesting questions in many pure disciplines: mathematics, computer science (computational geometry), statistics
- Interest from domain experts in biology, neuroscience, computer vision, dynamical systems, sensor networks, ...
- Materials science, pattern formation
- Machine learning: small features matter
- Agent-based modeling (swarming)

Possible answer: Persistent homology measures both the local geometry and the global topology of a dataset.



Measures of Order for nearly hexagonal lattices by Francis Motta, Rachel Neville, Patrick Shipman, Daniel Pearson, and Mark Bradley, 2018.

Why is applied topology popular when few datasets have Klein bottles?

- Many datasets have clusters & flares (as in the diabetes example)
- Motivates interesting questions in many pure disciplines: mathematics, computer science (computational geometry), statistics
- Interest from domain experts in biology, neuroscience, computer vision, dynamical systems, sensor networks, ...
- Materials science, pattern formation
- Machine learning: small features matter
- Agent-based modeling (swarming)

Possible answer: Persistent homology measures both the local geometry and the global topology of a dataset.



Answer: (from left) $r = 1.75, 2, 1.75, 2, 2.$

Why is applied topology popular when few datasets have Klein bottles?

- Many datasets have clusters & flares (as in the diabetes example)
- Motivates interesting questions in many pure disciplines: mathematics, computer science (computational geometry), statistics
- Interest from domain experts in biology, neuroscience, computer vision, dynamical systems, sensor networks, ...
- Materials science, pattern formation
- Machine learning: small features matter
- Agent-based modeling (swarming)

Possible answer: Persistent homology measures both the local geometry and the global topology of a dataset.



Agent-Based Modeling

Collective phenomenon, self-organization

Dutch Starling murmuration
filmed by Roald van Stijn

https://www.youtube.com/watch?v=YjDYE5CUb7Q

Takens' Theorem

Roughly speaking: Let M be a d-dimensional compact manifold, let $\phi\colon M \to M$ be a flow, and let $f\colon M \to \mathbb{R}$ be a measurement. Then *generically*,

$$m \mapsto \big(f(m), f(\phi(m), f(\phi^2(m), \ldots, f(\phi^{2d}(m))\big)$$

is an embedding $M \hookrightarrow \mathbb{R}^{2d+1}$.



Detecting strange attractors in turbulence by Floris Takens, 1982

Conley index theory



Conclusions for Part I

- Datasets have shape, which are reflective of patterns within.
- Persistent homology is a way to measure some of the local geometry and global topology of a dataset.



Evasion Paths in Mobile Sensor Networks



Topology applied to sensor networks

- Sensors move in a ball-shaped domain $B \subset \mathbb{R}^d$ over time interval $I = [0, 1]$. Fixed sensors cover the boundary.
- Measure only the Čech complex.
- Is there an evasion path?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Sensors move in a ball-shaped domain $B \subset \mathbb{R}^d$ over time interval $I = [0, 1]$. Fixed sensors cover the boundary.
- Measure only the Čech complex.
- Is there an evasion path?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Sensors move in a ball-shaped domain $B \subset \mathbb{R}^d$ over time interval $I = [0, 1]$. Fixed sensors cover the boundary.
- Measure only the Čech complex.
- Is there an evasion path?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Sensors move in a ball-shaped domain $B \subset \mathbb{R}^d$ over time interval $I = [0, 1]$. Fixed sensors cover the boundary.
- Measure only the Čech complex.
- Is there an evasion path?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Sensors move in a ball-shaped domain $B \subset \mathbb{R}^d$ over time interval $I = [0, 1]$. Fixed sensors cover the boundary.
- Measure only the Čech complex.
- Is there an evasion path?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Let $X \subset B \times I$ be the covered region.
- An *evasion path* is a time-preserving map from I to the uncovered region.



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Let $X \subset B \times I$ be the covered region.
- An *evasion path* is a time-preserving map from I to the uncovered region.
- <u>Evasion Problem.</u> Using the time-varying Čech complex, can we determine if an evasion path exists?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Let $X \subset B \times I$ be the covered region.
- An *evasion path* is a time-preserving map from I to the uncovered region.
- <u>Evasion Problem.</u> Using the time-varying Čech complex, can we determine if an evasion path exists?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- Let $X \subset B \times I$ be the covered region.
- An *evasion path* is a time-preserving map from I to the uncovered region.
- <u>Evasion Problem.</u> Using the time-varying Čech complex, can we determine if an evasion path exists?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- <u>Theorem (de Silva, Ghrist).</u>



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- <u>Theorem (de Silva, Ghrist).</u>



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- <u>Theorem (de Silva, Ghrist).</u>



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- <u>Theorem (de Silva, Ghrist).</u>
  If there is an $\alpha \in H_d(SC, \partial B \times I)$ with
  $0 \neq \partial\alpha \in H_{d-1}(\partial B \times I)$, then no evasion path exists.



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- <u>Theorem (de Silva, Ghrist).</u>
  If there is an $\alpha \in H_d(SC, \partial B \times I)$ with
  $0 \neq \partial\alpha \in H_{d-1}(\partial B \times I)$, then no evasion path exists.
- Coordinate-free.
- Not sharp. Can it be sharpened?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Topology applied to sensor networks

- <u>Theorem (de Silva, Ghrist).</u>
  If there is an $\alpha \in H_d(SC, \partial B \times I)$ with
  $0 \neq \partial\alpha \in H_{d-1}(\partial B \times I)$, then no evasion path exists.
- Coordinate-free.
- Not sharp. Can it be sharpened?



Coordinate-free Coverage in Sensor Networks with Controlled Boundaries via Homology by V. de Silva and R. Ghrist

Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



Zigzag Persistence by G. Carlsson and V. de Silva

Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



- <u>Theorem.</u>

  If there is an evasion path then there is a full-length bar.

Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



- <u>Theorem.</u>

  If there is an evasion path then there is a full-length bar.

Zigzag persistent homology

Form zigzag module for $X \longrightarrow I$ with $(d-1)$–dimensional homology.



- <u>Theorem.</u>

  If there is an evasion path then there is a full-length bar.

- Streaming computation.

Dependence on embedding $X \hookrightarrow B \times I$

- The time-varying Čech complex of X does not determine if an evasion path exists!



Dependence on embedding $X \hookrightarrow B \times I$

- The time-varying Čech complex of X does not determine if an evasion path exists!



Dependence on embedding $X \hookrightarrow B \times I$

- The time-varying Čech complex of X does not determine if an evasion path exists!



Dependence on embedding $X \hookrightarrow B \times I$

- The time-varying Čech complex of X does not determine if an evasion path exists!



Dependence on embedding $X \hookrightarrow B \times I$

- The time-varying Čech complex of X does not determine if an evasion path exists!



Dependence on embedding $X \hookrightarrow B \times I$



Dependence on embedding $X \hookrightarrow B \times I$



- The two covered regions are “topologically indistinguishable in a time-preserving way”, but the uncovered regions are not!

Zigzag persistence

- <u>Caution 2.9 of *Zigzag Persistence*.</u> Not every submodule isomorphic to an interval corresponds to a summand.





Zigzag persistence

- <u>Caution 2.9 of *Zigzag Persistence*.</u> Not every submodule isomorphic to an interval corresponds to a summand.





Dependence on embedding $X \hookrightarrow B \times I$



- The two covered regions are “topologically indistinguishable in a time-preserving way”, but the uncovered regions are not!

Conclusions for Part II

- There is a streaming one-sided criterion for the evasion problem using zigzag persistence.
- Cech complex insufficient.
  Alpha complex with rotation information suffices.
  What about the Čech complex with rotation information?



Where can I find resources if I am interested in applied topology?

- You may be interested in the Applied Algebraic Topology Research Network. Become a member to receive email invites to the online research seminars. Recorded talks are available at the YouTube Channel. There is also a forum.
- Another source of applied topology news is appliedtopology.org.
- A second online research seminar is GEOTOP-A: Applications of Geometry and Topology.
- Mailing lists with announcements in applied topology include WinCompTop and ALGTOP-L.

https://www.math.colostate.edu/~adams/advising



Thank you!