

Computing Persistent Homology*

Afra Zomorodian¹ and Gunnar Carlsson²

¹Department of Computer Science, Stanford University,
Stanford, CA 94305, USA
afra@cs.stanford.edu

²Department of Mathematics, Stanford University,
Stanford, CA 94305, USA
gunnar@math.stanford.edu

Abstract. We show that the persistent homology of a filtered d -dimensional simplicial complex is simply the standard homology of a particular graded module over a polynomial ring. Our analysis establishes the existence of a simple description of persistent homology groups over arbitrary fields. It also enables us to derive a natural algorithm for computing persistent homology of spaces in arbitrary dimension over any field. This result generalizes and extends the previously known algorithm that was restricted to subcomplexes of S^3 and \mathbb{Z}_2 coefficients. Finally, our study implies the lack of a simple classification over non-fields. Instead, we give an algorithm for computing individual persistent homology groups over an arbitrary principal ideal domain in any dimension.

1. Introduction



In this paper we study the homology of a filtered d -dimensional simplicial complex K , allowing an arbitrary principal ideal domain D as the ground ring of coefficients. A filtered complex is an increasing sequence of simplicial complexes, as shown in Fig. 1. It determines an *inductive system* of homology groups, i.e., a family of Abelian groups $\{G_i\}_{i \geq 0}$ together with homomorphisms $G_i \rightarrow G_{i+1}$. If the homology is computed with field coefficients, we obtain an inductive system of vector spaces over the field. Each vector space is determined up to isomorphism by its dimension. In this paper we obtain a simple classification of an inductive system of vector spaces. Our classification is in

* Research by the first author was partially supported by NSF under Grants CCR-00-86013 and ITR-0086013. Research by the second author was partially supported by NSF under Grant DMS-0101364. Research by both authors was partially supported by NSF under Grant DMS-0138456.

Outline

- Persistent homology
- Equivalence with $F[t]$ -modules
- Decomposition theorem
- Computation
- Experiments


Persistent homology



Apply i -dimensional homology with coefficients in a field F to get

$$H_i(K_0) \rightarrow H_i(K_1) \rightarrow H_i(K_2) \rightarrow H_i(K_3) \rightarrow H_i(K_4) \rightarrow H_i(K_5)$$

Equivalence



A persistence module

$$\mathbb{V}: V_0 \xrightarrow{f_0} V_1 \xrightarrow{f_1} V_2 \xrightarrow{f_2} V_3 \xrightarrow{f_3} V_4 \xrightarrow{f_4} V_5$$

can be thought of as a graded $F[t]$ -module
 with elements $(v_0, v_1, v_2, v_3, v_4, v_5) \in V_0 \oplus \dots \oplus V_5$
 and $F[t]$ action given by:


$$3 \cdot (v_0, \dots, v_5) = (3v_0, \dots, 3v_5)$$

$$t \cdot (v_0, \dots, v_5) = (0, f_0(v_0), f_1(v_1), f_2(v_2), f_3(v_3), f_4(v_4))$$

$$t^2 \cdot (v_0, \dots, v_5) = (0, 0, f_1 f_0(v_0), f_2 f_1(v_1), f_3 f_2(v_2), f_4 f_3(v_3))$$

$$(3 + t^2) \cdot (v_0, \dots, v_5) = (3v_0, 3v_1, 3v_2 + f_1 f_0(v_0), \dots)$$

Computation




$$M_1 = \left[\begin{array}{c|ccccc} & ab & bc & cd & ad & ac \\ \hline d & 0 & 0 & t & t & 0 \\ c & 0 & 1 & t & 0 & t^2 \\ b & t & t & 0 & 0 & 0 \\ a & t & 0 & 0 & t^2 & t^3 \end{array} \right]$$

In our example, we have


$$\tilde{M}_1 = \left[\begin{array}{c|ccccc} & cd & bc & ab & z_1 & z_2 \\ \hline d & \boxed{t} & 0 & 0 & 0 & 0 \\ c & t & \boxed{1} & 0 & 0 & 0 \\ b & 0 & t & \boxed{t} & 0 & 0 \\ a & 0 & 0 & t & 0 & 0 \end{array} \right], \quad (8)$$

where $z_1 = ad - cd - t \cdot bc - t \cdot ab$ and $z_2 = ac - t^2 \cdot bc - t^2 \cdot ab$ form a homogeneous basis for Z_1 .

Sparse computation



Experiments



Klein bottle
Electrostatic charge
Jet engine

	$ K $	Length	Filtration (s)	Persistence (s)
K	12,000	1,020	0.03	< 0.01
E	529,225	3,013	3.17	5.00
J	3,029,383	256	24.13	50.23

