

Math 474 HW #5

Due 2:00 PM Friday, Oct. 4

1. (Shifrin Problem 1.3.9) Let $\alpha : [0, L] \rightarrow \mathbb{R}^3$ be parametrized by arclength. A *unit normal vector field* X is a vector-valued function with $X(0) = X(L)$, $X(s) \cdot T(s) = 0$, and $\|X(s)\| = 1$ for all s . Define the *twist* of X to be

$$\text{tw}(\alpha, X) = \frac{1}{2\pi} \int_0^L X'(s) \cdot (T(s) \times X(s)) \, ds.$$

- (a) Show that if X and Y are two unit normal fields on α , then $\text{tw}(\alpha, X)$ and $\text{tw}(\alpha, Y)$ differ by an integer.
- (b) From part (a), the fractional part of $\text{tw}(\alpha, X)$ is independent of X , so depends only on α . We call this the *total twist* of α . Prove that the total twist of α equals the fractional part of $\frac{1}{2\pi} \int_0^L \tau \, ds$.
- (c) Prove that if a closed curve lies on a sphere, then its total twist is 0.

2. For a *piecewise-smooth* closed curve in \mathbb{R}^3 , we can define the total curvature of the curve to be

$$\int \kappa(s) \, ds + \sum_{i=1}^n \theta_i,$$

where the θ_i are the exterior angles at the corners of the curve (see Shifrin's problem 1.3.12 for a precise definition of the exterior angle and a picture).

Generalize Fenchel's Theorem to show that the total curvature of a piecewise-smooth curve is $\geq 2\pi$, with equality if and only if the curve is planar and convex.

(*Hint:* Think about how to extend the definition of the tangent indicatrix to piecewise-smooth curves.)