

Pillow Degenerations of $K3$ Surfaces

C. Ciliberto, R. Miranda, and M. Teicher

April 16, 2001

1 Introduction

In this article we construct a specific projective degeneration of $K3$ surfaces of degree $2g - 2$ in \mathbb{P}^g to a union of $2g - 2$ planes, which meet in such a way that the combinatorics of the configuration of planes is a triangulation of the 2-sphere. Abstractly, such degenerations are said to be Type III degenerations of $K3$ surfaces, see [5], [10], [3]. Although the birational geometry of such degenerations is fairly well understood, the study of projective degenerations is not nearly as completely developed.

In [1], projective degenerations of $K3$ surfaces to unions of planes were constructed, in which the general member was embedded by a primitive line bundle. The application featured there was a computation of the rank of the Wahl map for the general hyperplane section curve on the $K3$ surface.

In this article we construct degenerations for which the general member is embedded by a *multiple* of the primitive line bundle class. The construction depends on two parameters, and we intend in follow-up work to use these degenerations to compute braid monodromy for Galois coverings, in the style of [7] and [8]. We hope that the freedom afforded by the additional discrete parameters in the construction will yield interesting phenomena related to fundamental groups.

The specific degenerations which we construct can be viewed as two rectangular arrays of planes, joined along their boundary; for this reason we have given them the name “pillow” degenerations. They are described in Section 3. Following that, in Section 4, we study the degeneration of the general branch curve (for a general projection of the surfaces to a plane) to a union of lines (which is the “branch curve” for the union of planes). In particular when the general branch curve is a plane curve having only nodes and cusps as singularities, we describe the degeneration of the nodes and the cusps to the configuration of the union of lines. This is critical information in the application to the computation of the braid monodromy.

We are not aware of a modern reference for the statement that the general branch curve for a linear projection of a surface to a plane has only nodes and cusps as singularities. In this article we will operate under the assumption that this “folklore” statement is true and proceed. The reader may wish to consult

[6] for further information. We have included a short section at the beginning of the article deriving the characters of a general branch curve (degree, number of nodes and cusps) for the convenience of the reader, under this assumption.

The authors are grateful to the NATO Scientific Affairs Division, the Ministry of Science of Israel, the Emmy Noether Research Institute of Mathematics at Bar-Ilan University, and EAGER (the European Union Research Network in Algebraic Geometry), for financial support for the Workshop in Eilat at which this article was completed.

2 Characters of a General Branch Curve

Here we briefly develop the formulas for the degree and number of nodes and cusps on a general branch curve B for a general projection of a smooth surface $S \subset \mathbb{P}^N$ to a general plane \mathbb{P}^2 , assuming that these are the only singularities. These formulas are not new, see for example [2], [4], but these standard references do much more, in either outdated notation or with much more advanced techniques, than are necessary for this more modest computation. Hence we thought it useful to include it here for completeness and for the convenience of the reader. The reader may also want to consult [6], [7], [9], and [11] for additional insight.

Denote by $\pi : S \rightarrow \mathbb{P}^2$ such a general projection. Let K and H be the canonical and hyperplane classes of S respectively. Let d be the degree of S and $g(H)$ the genus of a smooth hyperplane divisor. The intersection numbers KH and H^2 are related to d and g by

$$d = H^2 \quad \text{and} \quad 2g(H) - 2 = H^2 + KH. \quad (2.1)$$

The degree of the finite map π is equal to the degree d of the surface S . The degree b of the branch curve may be easily computed by noting that the pull-back of a line in \mathbb{P}^2 is a hyperplane divisor; hence the Hurwitz formula gives

$$2g(H) - 2 = d(-2) + \deg(B)$$

from which it follows that

$$b = \deg(B) = 2d + 2g(H) - 2 = 3d + KH. \quad (2.2)$$

Let $R \subset S$ denote the ramification curve, and denote by R_0 the residual curve (equal set-theoretically to the closure of $\pi^{-1}(B) - R$). R is a smooth curve, and the mapping π , restricted to R , is a desingularization of B .

Suppose that B has n nodes and k cusps and no other singularities. Over a general smooth point of B , the map π has $d - 1$ preimages, one on the ramification curve and $d - 2$ on the residual curve. Over each node of B , the ramification curve R has two smooth branches, and over each cusp, R has one smooth branch. Over a node of B , the residual curve R_0 meets R once transversally at each branch of R , and otherwise has $d - 4$ nodes of its own. Over a cusp of B , the residual curve R_0 meets R twice at the point of R lying over the cusp,

and is smooth there; it otherwise has $d - 3$ cusps of its own. In any case, over either a node or a cusp of B , there are only $d - 2$ preimages, instead of the $d - 1$ preimages over a general point of B . Therefore, computing Euler numbers, we see that

$$e(R \cup R_0) = e(\pi^{-1}(B)) = (d - 1)e(B) - (n + k). \quad (2.3)$$

The genus of the ramification curve R , being a desingularization of the branch curve B , is

$$g(R) = (b - 1)(b - 2)/2 - (n + k)$$

using Plücker's formulas. Its Euler number is therefore

$$e(R) = 2 - 2g(R) = 2(n + k) - b^2 + 3b.$$

Since R and B differ, topologically, only over the nodes, we see that the Euler number of B is

$$e(B) = e(R) - n = n + 2k - b^2 + 3b.$$

Letting $e(S)$ be the Euler number of the surface S , we see that

$$\begin{aligned} e(S) &= d[e(\mathbb{P}^2) - e(B)] + e(R \cup R_0) \\ &= 3d - de(B) + (d - 1)e(B) - (n + k) \quad \text{using (2.3)} \\ &= 3d - e(B) - n - k \\ &= 3d - [n + 2k - b^2 + 3b] - n - k \\ &= 3d + b^2 - 3b - 2n - 3k, \end{aligned}$$

so that

$$2n + 3k = 3d + b^2 - 3b - e(S). \quad (2.4)$$

Pulling back 2-forms via π , we have the standard formula that

$$K_S = \pi^*(K_{\mathbb{P}^2}) + R = -3H + R$$

and since $bH = \pi^*(B)$, we see that

$$2R + R_0 = \pi^*(B) = bH,$$

so that, as classes on S ,

$$R = K + 3H \quad \text{and} \quad R_0 = bH - 2R = -2K + (b - 6)H.$$

Since R and R_0 meet transversally at each of the two points of R over a node, and meet to order two at the point of R lying over a cusp, we see that $R \cdot R_0 =$

$2(n+k)$. Therefore $2n+2k = R \cdot R_0 = (K+3H)(-2K+(b-6)H)$; multiplying this out gives

$$2n+2k = -2K^2 + (b-12)KH + (3b-18)H^2. \quad (2.5)$$

Subtracting (2.5) from (2.4) gives

$$k = 3d + b^2 - 3b - e(S) + 2K^2 - (b-12)KH - (3b-18)d$$

and then one can solve either expression for the number of nodes. Simplifying the expressions somewhat leads to the following.

Proposition 2.6 *Let S be a smooth surface of degree d in \mathbb{P}^N , and let $\pi : S \rightarrow \mathbb{P}^2$ be a general projection. Let K and H be the canonical and hyperplane classes of S , respectively. Let B be the branch curve of the projection π , which is assumed to be a plane curve of degree b with n nodes, k cusps, and no other singularities. Then:*

- (a) $\deg(\pi) = \deg(S) = d = H^2$.
- (b) The degree of the branch curve B is $b = 3d + KH$.
- (c) The number of nodes of the branch curve B is

$$n = -3K^2 + e(S) + 24d + \frac{b^2}{2} - 15b.$$

- (d) The number of cusps of the branch curve B is

$$k = 2K^2 - e(S) - 15d + 9b.$$

- (e) Under a general projection of the branch curve B to a line, the number t of turning points (simple branch points) is

$$t = e(S) - 3d + 2b.$$

The last computation of turning points is obtained from the Hurwitz formula, applied to the ramification curve R , noting that there are simple branch points for such a projection at the points of R lying over the cusps of B also.

Example 2.7 (Veronese Surfaces) *Let S be the r^{th} Veronese image of \mathbb{P}^2 . In this case, if L denotes the line class of S , then $L^2 = 1$, $K = -3L$, and $H = rL$; hence $K^2 = 9$, $KH = -3r$, and $d = H^2 = r^2$. The Euler number $e(S) = 3$. Therefore*

$$\begin{aligned} b &= 3r(r-1); & n &= 3(r-1)(r-2)(3r^2+3r-8)/2; \\ k &= 3(r-1)(4r-5); & t &= 3(r-1)^2. \end{aligned}$$

Example 2.8 (Rational Normal Scrolls) Let S be a rational normal scroll, e.g. $\mathbb{P}^1 \times \mathbb{P}^1$ embedded by the complete linear system H of type $(1, r)$. The canonical class is of type $(-2, -2)$, so that $K^2 = 8$, $KH = -2r - 2$, and $d = H^2 = 2r$. The Euler number $e(S) = 4$. Therefore

$$b = 4r - 2; \quad n = 4(r - 1)(2r - 3); \quad k = 6r - 6; \quad t = 2r.$$

Example 2.9 (Del Pezzo Surfaces) Let S be a Del Pezzo Surface of degree d in \mathbb{P}^d , for $3 \leq d \leq 9$. Then S is isomorphic to the plane blown up at $9 - d$ points; if L denotes the class of a line, and E the sum of the classes of the $9 - d$ exceptional divisors, then $L^2 = 1$, $LE = 0$, and $E^2 = d - 9$; also $K = -3L + E$, and $H = -K$, so that $K^2 = H^2 = d$, and $KH = -d$. The Euler number $e(S) = 12 - d$. Therefore

$$b = 2d; \quad n = 2(d - 2)(d - 3) = 2d^2 - 10d + 12; \quad k = 6(d - 2); \quad t = 12.$$

Example 2.10 (K3 Surfaces) Let S be a K3 surface of degree $d = 2g - 2$ in \mathbb{P}^g . The canonical class is trivial, so that $K^2 = KH = 0$. The Euler number $e(S) = 24$. Therefore

$$b = 6g - 6; \quad n = 6(g - 2)(3g - 7) = 18g^2 - 78g + 84; \\ k = 24(g - 2); \quad t = 6g + 18.$$

3 Construction of the Pillow Degeneration


A non-hyperelliptic K3 surface of genus $g \geq 3$ can be embedded by the sections of a very ample line bundle as a smooth surface of degree $2g - 2$ in \mathbb{P}^g . When the line bundle generates the Picard group of the K3 surface, the embedded K3 surface can be degenerated to a union of $2g - 2$ planes in a variety of ways (see for example [1]). In this section we will describe a degeneration, which we call the *pillow* degeneration, which smooths to a K3 surface whose Picard group is generated by a sub-multiple of the hyperplane class.

Fix two integers a and b at least two; set $g = 2ab + 1$. The number of planes in the pillow degeneration is then $2g - 2 = 4ab$.

This projective space has $g + 1 = 2ab + 2$ coordinate points, and each of the $4ab$ planes is obtained as the span of three of these. The sets of three are indicated in Figure 1, which describes the bottom part of the "pillow" and the top part of the "pillow", which are identified along the boundaries of the two configurations. The reader will see that the boundary is a cycle of $2a + 2b$ lines.

Note that no three of the planes meet in a line. Also note that the set of bottom planes lies in a projective space of dimension $ab + a + b$, as do the set of top planes; these two projective spaces meet exactly along the span of the $2a + 2b$ boundary points, which has dimension $2a + 2b - 1$. Finally note that the four corner points of the pillow degeneration (labeled 1, $a + 1$, $a + b + 1$, and $a + 2b + 1$) are each contained in three distinct planes, while all other points are each contained in six planes. This property, that the number of lines and planes incident on each of the points is bounded, is important for the later

Figure 1: Configuration of Planes, Top and Bottom



computations, and is a feature of the pillow degeneration that is not available in other previous degenerations.

We will call such a configuration of planes a *pillow of bidegree* (a, b) .

Theorem 3.1 *For any a and b at least 2, the pillow of bidegree (a, b) is a degeneration of a smooth K3 surface of degree $4ab$ in a projective space of dimension $g = 2ab + 1$. If $c = \text{g.c.d}(a, b)$, then the general such K3 surface will have Picard group generated by a line bundle L such that cL is the hyperplane bundle.*

The proof of the Theorem will be made in three steps. First we will exhibit a degeneration of the K3 surface to a union of two rational surfaces, each isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$, embedded via the sections of the linear system of bidegree (a, b) . These two rational surfaces will meet along an elliptic normal curve which is anticanonical in each. Secondly we will simultaneously degenerate each rational surface to a union of ab quadrics, resulting in a total of $2ab$ quadrics. Finally we will degenerate each quadric to a union of two planes.

Proof: (Step One:) Note that the sections of the linear system of bidegree (a, b) embed $\mathbb{P}^1 \times \mathbb{P}^1$ as a surface in a projective space of dimension $ab + a + b$. Choose an anticanonical divisor (of bidegree $(2, 2)$) which is a smooth elliptic curve; it is mapped by the above embedding to an elliptic normal curve in a subspace of dimension $2a + 2b - 1$.

In our original ambient space of dimension $2ab + 1$, choose two subspaces of dimension $ab + a + b$ which meet along a subspace of dimension $2a + 2b - 1$. Make the above identical construction of the $\mathbb{P}^1 \times \mathbb{P}^1$ in each of the two subspaces, taking care to have the two elliptic normal curves identified in the intersection subspace.

This union R of the two rational surfaces is a degeneration of an embedded $K3$ surface, by an argument identical to that presented for Theorems 1 and 2 of [1], which we will not repeat in detail here. Briefly, one first checks via standard calculations that $H^0(N_R)$ has dimension $g^2 + 2g + 19$ and that $H^1(N_R) = H^2(N_R) = 0$. Secondly, the natural map from $H^0(N_R)$ to $H^0(T^1)$ is seen to be surjective. This is sufficient to prove that R represents a smooth point of its Hilbert scheme, whose general member is a smooth $K3$ surface of degree $2g - 2$.

(Step Two:) The second step can be achieved as in [7] by observing that each $\mathbb{P}^1 \times \mathbb{P}^1$ can be degenerated to a union of ab quadrics by degenerating the first coordinate \mathbb{P}^1 to a chain of a lines, and the second coordinate \mathbb{P}^1 to a chain of b lines. In this degeneration the elliptic curve degenerates to a cycle of $2(a + b)$ lines. This degeneration is made simultaneously for each of the two $\mathbb{P}^1 \times \mathbb{P}^1$'s, resulting in a degeneration to a union of $2ab$ quadrics. This configuration of $2ab$ quadrics meet as in Figure 1, without the diagonal lines: if one removes the diagonal lines from Figure 1 we obtain $2ab$ rectangles, each indicating a quadric. Each of these quadrics meets the others along a cycle of four lines (two vertical and two horizontal).

(Step Three:) Finally degenerate each quadric to a union of two planes, as in Figure 1. These degenerations can be executed completely independently of course, and it is elementary to see that this can be done keeping the four lines along which any one of the quadrics meet the others fixed.

(Step Four:) Finally note that if $c \neq 1$, the pillow degeneration of bidegree (a, b) is a degeneration of the c -uple embedding of the pillow degeneration of bidegree $(a/c, b/c)$. To see this, one uses the standard triangular degeneration of the Veronese embedding of the plane as described in [8].

The final point to check is that the general $K3$ surface in this 19-dimensional family has Picard group generated by $(1/c)H$, where H is the hyperplane class. Since we have a 19-dimensional family of $K3$ surfaces, the only question to be decided is which sub-multiple of the hyperplane system is the generator of the general Picard group. The maximum possible is the g.c.d c . Since the pillow is a c -uple Veronese, the hyperplane class is at least a c -fold multiple, and since it cannot be any more, this shows that the Picard group is generated by $(1/c)H$. This completes the proof of the Theorem.

Q.E.D

Note that in this degeneration, the horizontal and vertical lines appear first, and the diagonal lines appear second.

4 The Degeneration of the Branch Curve

We assume that we are in a general enough situation that for a generic projection of a $K3$ surface of degree g in \mathbb{P}^g to a plane, the branch curve is a curve of degree $6g - 6$, having $6(g-2)(3g-7)$ nodes and $24(g-2)$ cusps and no other singularities; these numbers were presented in Section 2. If one projects this branch curve onto a general line, the projection will have $6g + 18$ simple branch points.

It is our goal in this section to describe how these nodes, cusps, and branch points degenerate in a pillow degeneration.

Firstly, since the pillow degeneration consists entirely of planes, under a general projection each plane will map isomorphically onto the target plane. Therefore the degenerate branch curve is composed of the $3g - 3$ planar lines which are the images of the $3g - 3$ double lines of the pillow degeneration where two planes meet. Each of the $3g - 3$ planar lines have multiplicity two in the limit branch curve.

We see therefore that the general branch curve (of degree $6g - 6$) degenerates as a curve to the $3g - 3$ planar lines, each doubled. Our next task is to describe the degeneration of the nodes, cusps and branch points of the general branch curve. In any case it is clear that these distinguished points of the general branch curve can only go to points of the $3g - 3$ planar lines.

Secondly, it is elementary to compute that there are $(9/2)g^2 - (51/2)g + 39$ pairs of disjoint lines in the pillow degeneration. Each of these pairs of disjoint lines gives rise to an intersection of two planar line components of the limit branch curve. We refer to these points as *2-points* of the configuration of the $3g - 3$ planar lines.

In addition to these 2-points, we have exactly four *3-points*, corresponding to the projection of the four points in the pillow degeneration where 3 planes (and 3 double lines) meet. Finally we have $g - 3$ *6-points* corresponding to the projection of the $g - 3$ points in the pillow degeneration where 6 planes (and 6 double lines) meet. At any one of these n -points ($n = 2, 3$, or 6) exactly n of the $3g - 3$ planar lines meet; moreover at no other point of the plane do any of these lines meet.

In the degeneration of the general branch curve to this configuration of $3g - 3$ double lines, each of the nodes, cusps, and branch points can degenerate either to a 2-point, a 3-point, a 6-point, or a smooth point of one of the $3g - 3$ lines. With the above terminology, we can now describe how many nodes, cusps, and branch points degenerate to each of these types of points.

Theorem 4.1 *In the pillow degeneration of a K3 surface of degree g in \mathbb{P}^g , the nodes, cusps, and branch points of the general branch curve degenerate to the 2-points, 3-points, 6-points, and other smooth points of lines according to the following table:*

Object Type	Number	Branch Points	Nodes	Cusps
Lines	$3g - 3$	0	0	0
3-points	4	9	0	6
6-points	$g - 3$	6	24	24
2-points	$\frac{9}{2}g^2 - \frac{51}{2}g + 39$	0	4	0
Totals:		$6g + 18$	$18g^2 - 78g + 84$	$24(g - 2)$

In particular no node, cusp, or branch point degenerates to a smooth point of any of the $3g - 3$ double lines of the limit branch curve.

Proof: We first look at the row of the table for the 2-points. Since each of

the planar lines have multiplicity two in the branch curve, this crossing point actually is a limit of 4 nodes of the general branch curve (the 4 nodes appearing as the four intersection points of two pairs of lines). No cusp or branch point of the general branch curve has this crossing point as a limit in general, since these points are created by the projection of unrelated disjoint lines in the union of planes in \mathbb{P}^g .

We now turn our attention to the images of the multiple points of the pillow degeneration where n planes (and n double lines) meet at one point. We assume that $3 \leq n \leq 6$ in what follows. (In the pillow degeneration we have $n = 3$ or $n = 6$ only.) Under the generic planar projection, such points go to intersections of n of the corresponding planar lines. We will refer to these as *n-points* of the limit branch curve.

In order to analyze the number of nodes, cusps, and branch points of the general curve which go to these *n-points*, we make a local analysis near the multiple point of the union of planes. There are n planes incident to this multiple point, and they together span a \mathbb{P}^n . Locally this collection of n planes in \mathbb{P}^n smooths to a Del Pezzo surface of degree n . In a generic projection for such a Del Pezzo, the branch curve has degree $2n$, with $2(n-2)(n-3)$ nodes and $6n-12$ cusps; the number of simple branch points for this curve under generic projection to a line is 12.

The limit branch curve corresponding to the degeneration of the Del Pezzo to the union of n planes is a union of n lines concurrent at a point p , the images of the n lines through the multiple point.

A partial smoothing of the union of n planes may be obtained by taking two adjacent planes and smoothing them to a quadric surface. The corresponding smoothing of the limit branch curve smooths exactly one of the n lines to a conic, which is necessarily tangent to two adjacent lines. As the conic degenerates to the (double) line L , we see that no nodes of the general branch curve go to any point of L which is not p , and no cusps do either. The conic has two general branch points for a projection to a line, and one of these branch points goes to p and one does not.

This local analysis of this partial smoothing shows that in a complete smoothing to the Del Pezzo, no node can go to a point of any line except the concurrent point p , and neither can any cusp. Therefore all of the $2(n-2)(n-3)$ nodes degenerate to the concurrent point, and all of the $6n-12$ cusps do too. Moreover, of the 12 branch points for the general curve, all but n of them go to the concurrent point p . (The other n go to one on each line.)

In the cases $n = 3$ and $n = 6$ of interest in the pillow degeneration, the above analysis shows that arbitrarily close to a 3-point there are $9 = 12 - 3$ branch points, and no nodes and 6 cusps. Arbitrarily close to a 6-point there are $6 = 12 - 6$ branch points, and 24 nodes and 24 cusps. This gives the entries in the 3-point and 6-point rows of the table.

If we now total the number of branch points, nodes and cusps which degenerate to these multiple points, we obtain the values in the last row of the table. Since these are exactly the number of branch points, nodes, and cusps of the general curve, we must have accounted for all of the branch points, nodes, and

cusps already. In particular there are none left to degenerate to smooth points of the double lines.

This completes the proof of the Theorem.

Q.E.D

References

- [1] C. Ciliberto, A. Lopez, and R. Miranda: *Projective Degenerations of K3 Surfaces, Gaussian Maps, and Fano Threefolds*. Inv. math. **114** (1993), 641 – 667.
- [2] F. Enriques: *Le Superficie Algebriche*. Nicola Zanichelli Editore, Bologna 1949.
- [3] R. Friedman and D. Morrison: *The Birational Geometry of Degenerations: An Overview*, in: *The Birational Geometry of Degenerations*, R. Friedman and D. Morrison, Editors. Progress in Mathematics **29**, Birkhäuser Press 1983.
- [4] B. Iversen: *Critical points of an algebraic function*, Inv. math. **12** (1971), 210–224.
- [5] V. Kulikov: *Degenerations of K3 Surfaces and Enriques Surfaces*. Math. USSR Izvestija **11** (1977), 957–989.
- [6] V. Kulikov and Vik. Kulikov: *Generic coverings of the plane with A-D-E singularities*. Izvestiya RAN **64** (2000), 65–106
- [7] B. Moishezon and M. Teicher: *Simply Connected Algebraic Surfaces with Positive Index*, Inv. math. **89** (1987), 601–643.
- [8] B. Moishezon and M. Teicher: *Braid group techniques in complex geometry III: Projective degeneration of V_3* . In: *Classification of Algebraic Varieties*, Contemporary Mathematics **162**, AMS (1994), 313 – 332.
- [9] B. Moishezon, M. Teicher, and A. Robb: *On Galois covers of Hirzebruch surfaces* Math. Ann. **305** (1996), 493–539.
- [10] U. Persson and H. Pinkham: *Degeneration of surfaces with trivial canonical divisor*, Annals of Math. **113** (1981), 45–66.
- [11] M. Teicher and A. Robb: *Applications of Braid Group Techniques to the decomposition of moduli spaces, new examples*. Topology and its Applications **78** (1997), 143–151.